

TANZANIA TREE SEED AGENCY



HANDLING SEED OF *PELTOPHORUM PTEROCARPUM* (COPPER POD)

Seed Issue Note No. (61) 1996

Temporary storage

Seeds can be stored for 4 years at room temperature without significant loss of viability, if they are maintained in the sealed polythene bag. However, it is recommended to sow the seeds within one year of receipt.

Treatment before sowing

Pour boiling water over the seed (volume of seed to water should be 1:5), stir to facilitate soaking until the water-cools. Continue soaking for 12 hours.

Alternatively, seeds can be nipped at the end opposite the scar by nail cutter.

Sowing

A space about 1 cm should be left between the seeds. This will give a sowing density of 10,000 seeds per m² of seedbed corresponding to approx. 0.7 kg of seed per m² of seedbed. Seeds should be covered with 0.5 cm of sand.

Germination

At 4th day after sowing the root comes out. Germination reaches 40% after 7 days and 90% after 14 days.

One kg contains 14,000 seeds and may produce 12,000 seedlings under ideal conditions. However, in many cases only 10,000 seedlings per kg will be obtained.

Tanzania Tree Seed Agency
S.L.P. 373, Morogoro – Tanzania

Tel: 023 260 3192/3903

Fax: 023 260 3275

E-mail: ttsa@morogoro.net