



**HANDLING SEED OF *VITEX MOMBASSAE*
(SMELLY BERRY)**

Seed Issue Note No. (64) 1996

Temporary storage

Properly dried "nuts" can be stored at room temperature for about one year.

Treatment before sowing

Soak the nuts in cold water for 12 hrs and sun dried for 12 hours alternatively for two days.


Sowing

A space about 1.5 cm should be left between the nuts. This will give sowing density of 4,500 nuts per m² of seedbed corresponding to approx 2 kg of nuts per m² seed bed. Seeds should be covered with 0.5 cm of sand.

Germination

The germination process takes long time. At the 3rd week after sowing the root comes out (see drawing). One "nut" may produce more than one seedling. The germination reaches 30% after 5 weeks and 60% after 8 weeks.

One kg contains 2,500 nuts and may produce 1,500 seedlings under ideal conditions. In many cases, only 1,000 seedlings per kg of nuts will be obtained. Transplanting into pots can take place when the seedling has appearance shown at week 5 in the drawing.



Germination stages in seed of *Vitex mombassae*

**Tanzania Tree Seed Agency
S.L.P. 373, Morogoro – Tanzania**

Tel: 023 260 3192/3903

Fax: 023 260 3275

E-mail: ttsa@morogoro.net